

Heterotic Supersymmetry

Hans Peter Nilles

Physikalisches Institut

Universität Bonn



Messages from the heterotic string

Localization properties of quarks, leptons and Higgses

- Higgs bosons and top-quark in the “bulk” lead to large top-quark Yukawa coupling
- first 2 families localized (exhibiting family symmetries)

Messages from the heterotic string

Localization properties of quarks, leptons and Higgses

- Higgs bosons and top-quark in the “bulk” lead to large top-quark Yukawa coupling
- first 2 families localized (exhibiting family symmetries)

Mirage scheme for SUSY breakdown

- compressed spectrum for gauginos
- reduced fine tuning
- hierarchy of soft terms

Remnants of $N=4$ SUSY from higher dimensions that might hide Susy at the LHC!

The Pattern

A specific pattern for the soft masses with a large gravitino mass in the multi-TeV range ($< O(30)\text{TeV}$)

(Krippendorff, Nilles, Ratz, Winkler, 2012)

- normal squarks and sleptons in multi-TeV range
- top squarks (\tilde{t}_L, \tilde{b}_L) and \tilde{t}_R in TeV range
(suppressed by $\log(M_{\text{Planck}}/m_{3/2}) \sim 4\pi^2$)
- A-parameters in TeV range
- gaugino masses in TeV range
- mirage pattern for gaugino masses
(compressed spectrum)

emerging from realistic models of the “MiniLandscape”.

Geography

Many properties of the models depend on the geography of extra dimensions, such as

- the **location** of quarks and leptons,
- the **relative location** of Higgs bosons,

Geography

Many properties of the models depend on the geography of extra dimensions, such as


- the **location** of quarks and leptons,
- the **relative location** of Higgs bosons,

but there is also a “localization” of gauge fields


- $E_8 \times E_8$ in the bulk
- smaller gauge groups on various branes

Observed 4-dimensional gauge group is common subgroup of the various localized gauge groups!

Calabi Yau Manifold



Orbifold



Localization

Quarks, Leptons and Higgs fields can be localized:

- in the Bulk ($d = 10$ **untwisted** sector)
- on 3-Branes ($d = 4$ twisted sector **fixed points**)
- on 5-Branes ($d = 6$ twisted sector **fixed tori**)

Localization

Quarks, Leptons and Higgs fields can be localized:


- in the Bulk ($d = 10$ **untwisted** sector)
- on 3-Branes ($d = 4$ twisted sector **fixed points**)
- on 5-Branes ($d = 6$ twisted sector **fixed tori**)

but there is also a “localization” of gauge fields

- $E_8 \times E_8$ in the bulk
- smaller gauge groups on various branes


Observed 4-dimensional gauge group is common subgroup of the various localized gauge groups!

Localized gauge symmetries



(Förste, HPN, Vaudrevange, Wingerter, 2004)

Standard Model Gauge Group



The MiniLandscape

- many models with the **exact spectrum of the MSSM** (absence of chiral exotics)

(Lebedev, HPN, Raby, Ramos-Sanchez, Ratz, Vaudrevange, Wingerter, 2007-2009)

- **family symmetries for the first two families**

- gauge- and (partial) Yukawa unification

(Raby, Wingerter, 2007)

- **large top quark Yukawa coupling**

- models with **R-parity** + solution to the **μ -problem**

(Lebedev, HPN, Raby, Ramos-Sanchez, Ratz, Vaudrevange, Wingerter, 2007)

- gaugino condensation and **mirage mediation**

(Löwen, HPN, 2008)

A Benchmark Model

At the orbifold point the gauge group is

$$SU(3) \times SU(2) \times U(1)^9 \times SU(4) \times SU(2)$$

- one $U(1)$ is anomalous
- there are singlets and vectorlike exotics
- decoupling of exotics and breakdown of gauge group has been verified
- remaining gauge group

$$SU(3) \times SU(2) \times U(1)_Y \times SU(4)_{\text{hidden}}$$


- for discussion of neutrinos and R-parity we keep also the $U(1)_{B-L}$ charges

Spectrum

#	irrep	label	#	irrep	label
3	$(\mathbf{3}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{(1/6, 1/3)}$	q_i	3	$(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(-2/3, -1/3)}$	\bar{u}_i
3	$(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(1, 1)}$	\bar{e}_i	8	$(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{(0, *)}$	m_i
3 + 1	$(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(1/3, -1/3)}$	\bar{d}_i	1	$(\mathbf{3}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(-1/3, 1/3)}$	d_i
3 + 1	$(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{(-1/2, -1)}$	ℓ_i	1	$(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{(1/2, 1)}$	$\bar{\ell}_i$
1	$(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{(-1/2, 0)}$	h_d	1	$(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{(1/2, 0)}$	h_u
6	$(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(1/3, 2/3)}$	$\bar{\delta}_i$	6	$(\mathbf{3}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(-1/3, -2/3)}$	δ_i
14	$(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(1/2, *)}$	s_i^+	14	$(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(-1/2, *)}$	s_i^-
16	$(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(0, 1)}$	\bar{n}_i	13	$(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(0, -1)}$	n_i
5	$(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{2})_{(0, 1)}$	$\bar{\eta}_i$	5	$(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{2})_{(0, -1)}$	η_i
10	$(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{2})_{(0, 0)}$	h_i	2	$(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{2})_{(0, 0)}$	y_i
6	$(\mathbf{1}, \mathbf{1}; \mathbf{4}, \mathbf{1})_{(0, *)}$	f_i	6	$(\mathbf{1}, \mathbf{1}; \bar{\mathbf{4}}, \mathbf{1})_{(0, *)}$	\bar{f}_i
2	$(\mathbf{1}, \mathbf{1}; \mathbf{4}, \mathbf{1})_{(-1/2, -1)}$	f_i^-	2	$(\mathbf{1}, \mathbf{1}; \bar{\mathbf{4}}, \mathbf{1})_{(1/2, 1)}$	\bar{f}_i^+
4	$(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(0, \pm 2)}$	χ_i	32	$(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(0, 0)}$	s_i^0
2	$(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(-1/6, 2/3)}$	\bar{v}_i	2	$(\mathbf{3}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(1/6, -2/3)}$	v_i

Unification

- Higgs doublets are in untwisted sector
- heavy top quark in untwisted sector
- μ -term protected by a discrete symmetry



- Minkowski vacuum before Susy breakdown (no AdS)
- solution to μ -problem (Casas, Munoz, 1993)
- first two families localized (smaller Yukawa couplings) exhibiting a discrete family symmetry

Emergent localization properties

The benchmark model illustrates some of the general properties of the MiniLandscape

- exactly two Higgs multiplets (no triplets)
- the top quark lives in the untwisted sector (as well as the Higgs multiplets)
- only one trilinear Yukawa coupling (all others suppressed)

Emergent localization properties


The benchmark model illustrates some of the general properties of the MiniLandscape

- exactly two Higgs multiplets (no triplets)
- the top quark lives in the untwisted sector (as well as the Higgs multiplets)
- only one trilinear Yukawa coupling (all others suppressed)

The fact that the top-quark has this unique property among all the quarks and leptons has important consequences for the phenomenological predictions including supersymmetry breakdown.

(Krippendorf, HPN, Ratz, Winkler, 2012)

Heterotic string: gaugino condensation



Gravitino mass $m_{3/2} = \Lambda^3 / M_{\text{Planck}}^2$ and $\Lambda \sim \exp(-\tau)$

(Lebedev, HPN, Raby, Ramos-Sanchez, Ratz, Vaudrevange, Wingerter, 2006)

Heterotic string

Fixing U- and T- moduli in a supersymmetric way

(Kappl, Petersen, Raby, Ratz, Vaudrevange, 2010; Anderson, Gray, Lukas, Ovrut, 2011)


- we remain with a run-away dilaton

But we need to adjust the vacuum energy

- matter field in untwisted sector
- “downlifting” mechanism can fix τ as well (no need for nonperturbative corrections to the Kähler potential)

(Löwen, HPN, 2008)

Downlift



(Löwen, HPN, 2008)

Mirage scheme

Fixing U- and T- moduli in a supersymmetric way

(Kappl et al., 2010; Anderson et al., 2011)

- we remain with a run-away dilaton

But we need to adjust the vacuum energy

- matter field in untwisted sector
- “downlifting” mechanism can fix τ as well (no need for nonperturbative corrections to the Kähler potential)
- a so-called mirage scheme with suppression factor $\log(m_{3/2}/M_{\text{Planck}})$

(Löwen, HPN, 2008)

Susy breakdown via down-lifting

In string theory we have (from **flux** and **gaugino condensate**)

$$W = \text{flux} - \exp(-X)$$

- modulus mediation suppressed (from down-lifting)

$$X \sim \log(M_{\text{Planck}}/m_{3/2}) \sim 4\pi^2$$

- radiative corrections become relevant (β function)

- Mixed mediation scheme: **Mirage Mediation (MMAM)**
with mirage pattern for gaugino masses:


$$m_{1/2} \sim m_{3/2}/4\pi^2$$

(Choi, Falkowski, Nilles, Olechowski, 2005)


first encountered in the framework of Type IIB theory

(Kachru, Kallosh, Linde, Trivedi, 2003)

Evolution of couplings



The Mirage Scale



(Lebedev, HPN, Ratz, 2005)

Reading the Gaugino Code

Mixed boundary conditions at the GUT scale characterized by the **parameter** α :
the ratio of modulus to anomaly mediation.


- $M_1 : M_2 : M_3 \simeq 1 : 2 : 6$ for $\alpha \simeq 0$
- $M_1 : M_2 : M_3 \simeq 1 : 1.3 : 2.5$ for $\alpha \simeq 1$
- $M_1 : M_2 : M_3 \simeq 1 : 1 : 1$ for $\alpha \simeq 2$
- $M_1 : M_2 : M_3 \simeq 3.3 : 1 : 9$ for $\alpha \simeq \infty$

The mirage scheme leads to

- LSP χ_1^0 predominantly Bino
- a “compact” (compressed) gaugino mass pattern.

(Choi, HPN, 2007; Löwen, HPN, 2009)

Gaugino Masses



Soft terms

So we have mirage suppression (compared to $m_{3/2}$) of

- gaugino masses (with compressed spectrum)
- A-parameters in the (few) TeV range.

Scalar masses are less protected

- heavy squarks and sleptons: $m_0 < O(30)\text{TeV}$

(Lebedev, Nilles, Ratz, 2006; Löwen, Nilles, 2008)

But, the top quark plays a special role

- as a result of gauge-Yukawa-unification

$$g_{\text{top}} \sim g_{\text{gauge}} \sim g_{\text{string}}$$

that explains the large value of the top-quark mass

Soft terms

While normal scalar masses are less protected

- this is not true for the top- and Higgs-multiplets
- they live in the untwisted sector (bulk)
- all other multiplets live twisted sectors (branes)

This can be understood as a remnant of

- extended supersymmetry in higher dimensions
- $N = 4$ supersymmetry from $N = 1$ in $D = 10$ via torus compactification
- Higgs und stops remain in the TeV-range

(Krippendorf, Nilles, Ratz, Winkler, 2012)

The overall pattern

This provides a specific pattern for the soft masses with a large gravitino mass in the multi-TeV range


- normal squarks and sleptons in Multi-TeV range
- top squarks (\tilde{t}_L, \tilde{b}_L) and \tilde{t}_R in TeV-range
(suppressed by $\log(M_{\text{Planck}}/m_{3/2}) \sim 4\pi^2$)
- A-parameters in TeV range
- gaugino masses in TeV range
- mirage pattern for gaugino masses
(compressed spectrum)
- heavy moduli (enhanced by $\log(M_{\text{Planck}}/m_{3/2})$
compared to the gravitino mass)

A Closer Look

A more detailed picture requires the analysis of specific models. Issues that have to be clarified:


- the appearance of tachyons,
 - partially inherited from anomaly mediation
 - two loop effects in the presence of heavy scalars
- the hierarchy between gauginos and sfermions.
- Can we satisfy all phenomenological constraints?
 - mass of Higgs, correct electroweak symmetry breakdown etc.
 - nature and abundance of WIMP-LSP.
- What is the LHC reach to test this scheme?

Benchmark model with a TeV gluino




Parameter scan for a gluino mass of 1 TeV. The coloured regions are excluded while the hatched region indicates the current reach of the LHC. The contours indicate the mass of the lightest stop.

Spectrum 1




Model with 4 TeV gluino



Parameter scan for a gluino mass of 4 TeV. The coloured regions are excluded while the hatched region indicates the current reach of the LHC. The contours indicate the mass of the lightest stop.

Spectrum of model with a 4 TeV gluino



Messages

- large gravitino mass (multi TeV-range)
- heavy moduli: $m_{3/2} \log(M_{\text{Planck}}/m_{3/2})$
- mirage pattern for gaugino masses rather robust
- sfermion masses are of order $m_{3/2}$
- the ratio between sfermion and gaugino masses, however, seems to be limited
- the heterotic string yields “Natural SUSY”. There is a reduced fine-tuning because of
 - the mirage pattern for gauginos,
 - and light stop masses
- and this is a severe challenge for LHC searches.

Comparison to other schemes

Mirage pattern for gaugino masses seems to be common for type II, G2MSSM and heterotic models

- **type IIB**

- all sfermions unprotected
- A-parameters in few TeV-range

- **G2MSSM**

- all sfermions unprotected
- A-parameters in multi TeV-range (e.g. $O(50)$ TeV)

but there are **no explicit models** to test a connection between Yukawa pattern and soft breaking terms.

The overall scale

There is no (reliable) prediction for the gravitino mass

- except for fine-tuning arguments
- “no lose” criterion (SSC with 20+20 TeV)
- does LHC satisfy this criterion?

The overall scale

There is no (reliable) prediction for the gravitino mass

- except for fine-tuning arguments
- “no lose” criterion (SSC with 20+20 TeV)
- does LHC satisfy this criterion?

Betting in the early 80's

- I bet that supersymmetry will be discovered before SSC gets into operation
- I bet that supersymmetry will have been forgotten before SSC gets into operation

Conclusions

Assume realistic top-quark Yukawa coupling

- this implies Gauge-Yukawa unification (trilinear top quark Yukawa coupling)
- realistic models require Higgs multiplets and top multiplets in untwisted sector (GaugeHiggsUnification)
- other fields tend to be localized at fixed points (tori) and exhibit family symmetries

Conclusions

Assume realistic top-quark Yukawa coupling

- this implies Gauge-Yukawa unification (trilinear top quark Yukawa coupling)
- realistic models require Higgs multiplets and top multiplets in untwisted sector (GaugeHiggsUnification)
- other fields tend to be localized at fixed points (tori) and exhibit family symmetries

Remnants of N=4 SUSY

- mirage mediation
- untwisted SUSY partners rather light while others heavy