

# The Heterotic MSSM

Hans Peter Nilles

Physikalisches Institut

Universität Bonn

Germany



# Questions

- What can we learn from strings for particle physics?
- Can we incorporate particle physics models within the framework of string theory?

# Questions

- What can we learn from strings for particle physics?
- Can we incorporate particle physics models within the framework of string theory?

## Recent progress:

- explicit model building towards the MSSM
  - Heterotic brane world
  - local grand unification
- moduli stabilization and Susy breakdown
  - gaugino condensation and uplifting
  - mirage mediation

# The road to the Standard Model

What do we want?

- gauge group  $SU(3) \times SU(2) \times U(1)$
- 3 families of quarks and leptons
- scalar Higgs doublet

# The road to the Standard Model

What do we want?

- gauge group  $SU(3) \times SU(2) \times U(1)$
- 3 families of quarks and leptons
- scalar Higgs doublet

But there might be more:

- supersymmetry (SM extended to MSSM)
- neutrino masses and mixings

as a hint for a large mass scale around  $10^{16}$  GeV

# Indirect evidence

Experimental findings suggest the existence of two new scales of physics beyond the standard model

$M_{\text{GUT}} \sim 10^{16} \text{ GeV}$  (and  $M_{\text{SUSY}} \sim 10^3 \text{ GeV}$ ):

# Indirect evidence

Experimental findings suggest the existence of two new scales of physics beyond the standard model

$M_{\text{GUT}} \sim 10^{16} \text{ GeV}$  (and  $M_{\text{SUSY}} \sim 10^3 \text{ GeV}$ ):

- **Neutrino-oscillations** and “See-Saw Mechanism”

$$m_\nu \sim M_W^2 / M_{\text{GUT}}$$

$$m_\nu \sim 10^{-3} \text{ eV for } M_W \sim 100 \text{ GeV,}$$

# Indirect evidence

Experimental findings suggest the existence of two new scales of physics beyond the standard model

$M_{\text{GUT}} \sim 10^{16} \text{ GeV}$  (and  $M_{\text{SUSY}} \sim 10^3 \text{ GeV}$ ):

- **Neutrino-oscillations** and “See-Saw Mechanism”


$$m_\nu \sim M_W^2 / M_{\text{GUT}}$$

$$m_\nu \sim 10^{-3} \text{ eV for } M_W \sim 100 \text{ GeV,}$$


- **Evolution of couplings constants** of the standard model towards higher energies.



# MSSM (supersymmetric)



# Standard Model



# Grand Unification

This leads to SUSY-GUTs with nice things like

- unified multiplets (e.g. spinors of  $SO(10)$ )
- gauge coupling unification
- Yukawa unification
- neutrino see-saw mechanism

# Grand Unification

This leads to SUSY-GUTs with nice things like

- unified multiplets (e.g. spinors of  $SO(10)$ )
- gauge coupling unification
- Yukawa unification
- neutrino see-saw mechanism

But there remain a few difficulties:

- breakdown of GUT group (large representations)
- doublet-triplet splitting problem (incomplete multiplets)
- proton stability (need for R-parity)

# String Theory

What do we get from string theory?

- supersymmetry
- extra spatial dimensions
- large unified gauge groups
- consistent theory of gravity

# String Theory

What do we get from string theory?


- supersymmetry
- extra spatial dimensions
- large unified gauge groups
- consistent theory of gravity

These are the building blocks for a **unified theory** of all the fundamental interactions.


But do they fit together, and if yes how?

**We need to understand the mechanism of compactification of the extra spatial dimensions**

# Calabi Yau Manifold



# Orbifold





# Localization

Quarks, Leptons and Higgs fields can be localized:

- in the Bulk ( $d = 10$  **untwisted** sector)
- on 3-Branes ( $d = 4$  twisted sector **fixed points**)
- on 5-Branes ( $d = 6$  twisted sector **fixed tori**)

# Localization

Quarks, Leptons and Higgs fields can be localized:


- in the Bulk ( $d = 10$  **untwisted** sector)
- on 3-Branes ( $d = 4$  twisted sector **fixed points**)
- on 5-Branes ( $d = 6$  twisted sector **fixed tori**)

but there is also a “localization” of gauge fields


- $E_8 \times E_8$  in the bulk
- smaller gauge groups on various branes

Observed 4-dimensional gauge group is common subgroup of the various localized gauge groups!

# Localized gauge symmetries



# Standard Model Gauge Group



# Local Grand Unification

In fact string theory gives us a variant of GUTs

- complete multiplets for fermion families
- split multiplets for gauge- and Higgs-bosons
- partial Yukawa unification

# Local Grand Unification

In fact string theory gives us a variant of GUTs

- complete multiplets for fermion families
- split multiplets for gauge- and Higgs-bosons
- partial Yukawa unification

Key properties of the theory depend on the **geography** of the fields in extra dimensions.

This geometrical set-up called **local GUTs**, can be realized in the framework of the “heterotic braneworld”.

(Förste, HPN, Vaudrevange, Wingerter, 2004; Buchmüller, Hamaguchi, Lebedev, Ratz, 2004)

# The Remnants of $SO(10)$

- $SO(10)$  is realized in the higher dimensional theory
- broken in  $d = 4$
- coexistence of complete and incomplete multiplets

# The Remnants of $SO(10)$

- $SO(10)$  is realized in the higher dimensional theory
- broken in  $d = 4$
- coexistence of complete and incomplete multiplets


Still there could be remnants of  $SO(10)$  symmetry

- 16 of  $SO(10)$  at some branes
- correct hypercharge normalization
- R-parity
- distinction between different families

that are very useful for realistic model building ...



# The “fertile patch”: $Z_6$ II orbifold



(Kobayashi, Raby, Zhang, 2004; Buchmüller, Hamaguchi, Lebedev, Ratz, 2004)

- provides fixed points and fixed tori
- allows  $SO(10)$  gauge group
- allows for localized 16-plets for 2 families
- $SO(10)$  broken via Wilson lines
- nontrivial hidden sector gauge group

# Selection Strategy

| criterion                                | $V^{\text{SO}(10),1}$ | $V^{\text{SO}(10),2}$ |
|--|-----------------------|-----------------------|
| ② models with 2 Wilson lines             | 22,000                | 7,800                 |
| ③ SM gauge group $\subset \text{SO}(10)$ | 3563                  | 1163                  |
| ④ 3 net families                         | 1170                  | 492                   |
| ⑤ gauge coupling unification             | 528                   | 234                   |
| ⑥ no chiral exotics                      | 128                   | 90                    |

(Lebedev, HPN, Raby, Ramos-Sanchez, Ratz, Vaudrevange, Wingerter, 2006)

# The road to the MSSM

This scenario leads to

- 200 models with the **exact spectrum of the MSSM** (absence of chiral exotics)
- **local grand unification** (by construction)
- gauge- and (partial) Yukawa unification

(Raby, Wingerter, 2007)

- examples of **neutrino see-saw mechanism**

(Buchmüller, Hamguchi, Lebedev, Ramos-Sanchez, Ratz, 2007)

- models with **R-parity** + solution to the  **$\mu$ -problem**

(Lebedev, HPN, Raby, Ramos-Sanchez, Ratz, Vaudrevange, Wingerter, 2007)

- hidden sector gaugino condensation

# A Benchmark Model

At the orbifold point the gauge group is

$$SU(3) \times SU(2) \times U(1)^9 \times SU(4) \times SU(2)$$

- one  $U(1)$  is anomalous
- there are singlets and vectorlike exotics
- decoupling of exotics and breakdown of gauge group has been verified
- remaining gauge group

$$SU(3) \times SU(2) \times U(1)_Y \times SU(4)_{\text{hidden}}$$


- for discussion of neutrinos and R-parity we keep also the  $U(1)_{B-L}$  charges

# Spectrum


| #     | irrep  | label            | #  | irrep   | label         |
|-------|--|------------------|----|---|---------------|
| 3     | $(\mathbf{3}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{(1/6, 1/3)}$        | $q_i$            | 3  | $(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(-2/3, -1/3)}$ | $\bar{u}_i$   |
| 3     | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(1, 1)}$            | $\bar{e}_i$      | 8  | $(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{(0, *)}$             | $m_i$         |
| 3 + 1 | $(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(1/3, -1/3)}$ | $\bar{d}_i$      | 1  | $(\mathbf{3}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(-1/3, 1/3)}$        | $d_i$         |
| 3 + 1 | $(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{(-1/2, -1)}$        | $l_i$            | 1  | $(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{(1/2, 1)}$           | $\bar{l}_i$   |
| 1     | $(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{(-1/2, 0)}$         | $h_d$            | 1  | $(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{(1/2, 0)}$           | $h_u$         |
| 6     | $(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(1/3, 2/3)}$  | $\bar{\delta}_i$ | 6  | $(\mathbf{3}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(-1/3, -2/3)}$       | $\delta_i$    |
| 14    | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(1/2, *)}$          | $s_i^+$          | 14 | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(-1/2, *)}$          | $s_i^-$       |
| 16    | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(0, 1)}$            | $\bar{n}_i$      | 13 | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(0, -1)}$            | $n_i$         |
| 5     | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{2})_{(0, 1)}$            | $\bar{\eta}_i$   | 5  | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{2})_{(0, -1)}$            | $\eta_i$      |
| 10    | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{2})_{(0, 0)}$            | $h_i$            | 2  | $(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{2})_{(0, 0)}$             | $y_i$         |
| 6     | $(\mathbf{1}, \mathbf{1}; \mathbf{4}, \mathbf{1})_{(0, *)}$            | $f_i$            | 6  | $(\mathbf{1}, \mathbf{1}; \bar{\mathbf{4}}, \mathbf{1})_{(0, *)}$       | $\bar{f}_i$   |
| 2     | $(\mathbf{1}, \mathbf{1}; \mathbf{4}, \mathbf{1})_{(-1/2, -1)}$        | $f_i^-$          | 2  | $(\mathbf{1}, \mathbf{1}; \bar{\mathbf{4}}, \mathbf{1})_{(1/2, 1)}$     | $\bar{f}_i^+$ |
| 4     | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(0, \pm 2)}$        | $\chi_i$         | 32 | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(0, 0)}$             | $s_i^0$       |
| 2     | $(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(-1/6, 2/3)}$ | $\bar{v}_i$      | 2  | $(\mathbf{3}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(1/6, -2/3)}$        | $v_i$         |

# Unification

- Higgs doublets are in untwisted (U3) sector
- trilinear coupling to the top-quark allowed
- threshold corrections (“on third torus”) allow unification at correct scale around  $10^{16}$  GeV



# Hidden Sector Susy Breakdown



Gravitino mass  $m_{3/2} = \Lambda^3 / M_{\text{Planck}}^2$  is in the **TeV range**  
for the hidden sector gauge group  $SU(4)$

(Lebedev, HPN, Raby, Ramos-Sanchez, Ratz, Vaudrevange, Wingerter, 2006)

# See-saw neutrino masses

The see-saw mechanism requires

- right handed neutrinos ( $Y = 0$  and  $B - L = \pm 1$ ),
- heavy Majorana neutrino masses  $M_{\text{Majorana}}$ ,
- Dirac neutrino masses  $M_{\text{Dirac}}$ .

The benchmark model has 49 right handed neutrinos:

- the left handed neutrino mass is  $m_\nu \sim M_{\text{Dirac}}^2 / M_{\text{eff}}$
- with  $M_{\text{eff}} < M_{\text{Majorana}}$  and depends on the number of right handed neutrinos.

(Buchmüller, Hamguchi, Lebedev, Ramos-Sanchez, Ratz, 2007;  
Lebedev, HPN, Raby, Ramos-Sanchez, Ratz, Vaudrevange, Wingerter, 2007)



# Spectrum

| #     | irrep  | label            | #  | irrep   | label         |
|-------|--|------------------|----|---|---------------|
| 3     | $(\mathbf{3}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{(1/6, 1/3)}$        | $q_i$            | 3  | $(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(-2/3, -1/3)}$ | $\bar{u}_i$   |
| 3     | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(1, 1)}$            | $\bar{e}_i$      | 8  | $(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{(0, *)}$             | $m_i$         |
| 3 + 1 | $(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(1/3, -1/3)}$ | $\bar{d}_i$      | 1  | $(\mathbf{3}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(-1/3, 1/3)}$        | $d_i$         |
| 3 + 1 | $(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{(-1/2, -1)}$        | $l_i$            | 1  | $(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{(1/2, 1)}$           | $\bar{l}_i$   |
| 1     | $(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{(-1/2, 0)}$         | $h_d$            | 1  | $(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{1})_{(1/2, 0)}$           | $h_u$         |
| 6     | $(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(1/3, 2/3)}$  | $\bar{\delta}_i$ | 6  | $(\mathbf{3}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(-1/3, -2/3)}$       | $\delta_i$    |
| 14    | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(1/2, *)}$          | $s_i^+$          | 14 | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(-1/2, *)}$          | $s_i^-$       |
| 16    | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(0, 1)}$            | $\bar{n}_i$      | 13 | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(0, -1)}$            | $n_i$         |
| 5     | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{2})_{(0, 1)}$            | $\bar{\eta}_i$   | 5  | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{2})_{(0, -1)}$            | $\eta_i$      |
| 10    | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{2})_{(0, 0)}$            | $h_i$            | 2  | $(\mathbf{1}, \mathbf{2}; \mathbf{1}, \mathbf{2})_{(0, 0)}$             | $y_i$         |
| 6     | $(\mathbf{1}, \mathbf{1}; \mathbf{4}, \mathbf{1})_{(0, *)}$            | $f_i$            | 6  | $(\mathbf{1}, \mathbf{1}; \bar{\mathbf{4}}, \mathbf{1})_{(0, *)}$       | $\bar{f}_i$   |
| 2     | $(\mathbf{1}, \mathbf{1}; \mathbf{4}, \mathbf{1})_{(-1/2, -1)}$        | $f_i^-$          | 2  | $(\mathbf{1}, \mathbf{1}; \bar{\mathbf{4}}, \mathbf{1})_{(1/2, 1)}$     | $\bar{f}_i^+$ |
| 4     | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(0, \pm 2)}$        | $\chi_i$         | 32 | $(\mathbf{1}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(0, 0)}$             | $s_i^0$       |
| 2     | $(\bar{\mathbf{3}}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(-1/6, 2/3)}$ | $\bar{v}_i$      | 2  | $(\mathbf{3}, \mathbf{1}; \mathbf{1}, \mathbf{1})_{(1/6, -2/3)}$        | $v_i$         |

# R-parity

- R-parity allows the **distinction** between Higgs bosons and sleptons
- $SO(10)$  **contains R-parity** as a discrete subgroup of  $U(1)_{B-L}$ .
- in conventional “**field theory GUTs**” one needs large representations to break  $U(1)_{B-L}$  (>126 dimensional)
- in **heterotic string** models one has more candidates for R-parity (and generalizations thereof)
- one just needs **singlets with an even  $B - L$  charge** that **break  $U(1)_{B-L}$  down to R-parity**

(Lebedev, HPN, Raby, Ramos-Sanchez, Ratz, Vaudrevange, Wingerter, 2007)

# Discrete Symmetries

There are numerous discrete symmetries

- from geometry
- and from stringy selection rules,
- both of abelian and nonabelian nature.

(Kobayashi, HPN, Plöger, Raby, Ratz, 2006)

Possible applications:

- (nonabelian) family symmetries
- Yukawa textures
- approximate global  $U(1)$  for a QCD axion

(Choi, Kim, Kim, 2006; Choi, HPN, Ramos-Sanchez, Vaudrevange, 2008)

# The $\mu$ problem

In general we have to worry about

- doublet-triplet splitting
- mass term for additional doublets
- the appearance of “naturally” light doublets

In the benchmark model we have

- only 2 doublets
- which are neutral under all selection rules
- if  $M(s_i)$  allowed in superpotential
- then  $M(s_i)H_uH_d$  is allowed as well

# The $\mu$ problem II

We have verified that (up to order 6 in the singlets)

- $F_i = 0$  implies automatically
- $M(s_i) = 0$  for all allowed terms  $M(s_i)$  in the superpotential  $W$


Therefore

- $W = 0$  in the supersymmetric (Minkowski) vacuum
- as well as  $\mu = \partial^2 W / \partial H_u \partial H_d = 0$ , while all the vectorlike exotics decouple
- with broken supersymmetry  $\mu \sim m_{3/2} \sim \langle W \rangle$

This solves the  $\mu$ -problem

(Casas, Munoz, 1993)

# Gaugino Condensation




Gravitino mass  $m_{3/2} = \Lambda^3 / M_{\text{Planck}}^2$  and  $\Lambda \sim \exp(-S)$


**We need to fix the dilaton!**

(Lebedev, HPN, Raby, Ramos-Sanchez, Ratz, Vaudrevange, Wingerter, 2006)

# Run-away potential



# Corrections to Kähler potential



(Casas, 1996; Barreiro, de Carlos, Copeland, 1998)



# Dilaton Domination?

This is known as the dilaton domination scenario,


- but there are problems to remove the vacuum energy.

One needs a “downlifting” mechanism:

- the analogue to the F-term “uplifting” in the type IIB case  
(Gomez-Reino, Scrucca, 2006; Lebedev, HPN, Ratz, 2006)
- “downlifting” mechanism fixes  $S$  as well (no need for nonperturbative corrections to the Kähler potential)  
(Löwen, HPN, 2008)
- mirage mediation for gaugino masses


# Sequestered sector “uplifting”

$N = 4$      $A = 4.9$      $C_0 = 0.73$




(Lebedev, HPN, Ratz, 2006; Löwen, HPN, 2008)

# Metastable “Minkowski” vacuum




(Löwen, HPN, 2008)


# Evolution of couplings



# The Mirage Scale




# Constraints on the mixing parameter



(Löwen, HPN, 2008)

# Constraints on the mixing parameter



(Löwen, HPN, 2008)

# Conclusion

String theory provides us with **new ideas for particle physics** model building, leading to concepts such as

- Local Grand Unification
- realistic MSSM candidates

**Geography of extra dimensions** plays a crucial role:

- localization of fields on branes,
- sequestered sectors and **mirage mediation**

LHC might help us to verify some of these ideas!