
Exercises on Elementary Particle Physics II

Prof. Dr. H.-P. Nilles

1. *D-term SUSY breaking - part II*

In this exercise we continue to discuss the details of D-term SUSY breaking (see Ex.7.2.). The U(1) gauge superfield is denoted by $V = (V_\mu, \lambda, D)$ and the left-chiral superfield by $\phi = (\varphi, \psi, F)$, having charge q .

(a) Case $q\xi < 0$.

From the shape of the scalar potential $V(\varphi)$, we see that the radial component of φ gets massive, while the angular component stays massless. Verify this by computation.

Since φ is charged, the vev of $\varphi\varphi^*$ breaks the U(1) gauge symmetry. Compute the mass of the gauge boson V_μ .

Is it possible to define a massive Dirac fermion? What is the relation between the Dirac mass and the mass of the gauge boson? Describe the SUSY-Higgs mechanism.

Hint: The mass of the gauge boson arises from the term $(D_\mu\varphi(x))(D^\mu\varphi(x))^*$. The massive Dirac fermion is composed of ψ and the gaugino λ .

(b) Case $q\xi > 0$.

Show that all particles except for φ stay massless. Compute the mass of φ .

2. *The Minimal Supersymmetric Standard Model - part I*

All standard model matter fields can be included in the following left-chiral superfields:

$$\begin{array}{lll} \text{quarks} & \mathbb{U}_i = (\mathbf{3}, \mathbf{2}, \frac{1}{6}) & \bar{\mathbb{U}}_i = (\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3}) \quad \bar{D}_i = (\bar{\mathbf{3}}, \mathbf{1}, \frac{1}{3}) \\ \text{leptons} & \mathbb{L}_i = (\mathbf{1}, \mathbf{2}, -\frac{1}{2}) & \bar{E}_i = (\mathbf{1}, \mathbf{1}, 1) \\ \text{higgs} & H = (\mathbf{1}, \mathbf{2}, -\frac{1}{2}) & \bar{H} = (\mathbf{1}, \mathbf{2}, \frac{1}{2}) \end{array}$$

The index $i = 1, 2, 3$ labels the three generations of quarks and leptons.

(a) How do the component fields (of e.g. \mathbb{L}_i) transform under gauge transformations? Why is the term $\mathbb{L}H\bar{E}$ gauge invariant?

- (b) Write the most general, gauge invariant cubic superpotential for these superfields.
- (c) Why do we need a second Higgs \bar{H} in the MSSM?
- (d) Identify the terms that conserve baryon and lepton number and those that do not. Verify that R-parity $R_p = (-1)^{3B+L+2s}$ forbids exactly those terms that violate baryon or lepton number.